

## Today

Continue Review: linear programming.  
 Simplex and Matching.  
 Taking Duals.

$$\begin{aligned} \max & 4x_1 + 2x_2 \\ & 2x_1 \leq 40 \\ & 3x_2 \leq 75 \\ & 3x_1 + 2x_2 \leq 100 \\ & x_1, x_2 \geq 0 \end{aligned}$$

Optimal point?  
 Try every point if we only had time!  
 How many points?  
 Real numbers?  
 Infinite. Uncountably infinite!

## Profit maximization.

Plant Carrots or Peas?

2\$ bushel of carrots. 4\$ for peas.

Carrots take 3 unit of water/bushel.  
 Peas take 2 units of water/bushel.

100 units of water.

Peas require 2 yards/bushel of sunny land.  
 Carrots require 1 yard/bushel of shadyland.


Garden has 60 yards of sunny land and 75 yards of shady land.

To pea or not to pea, that is the question!

A linear program.

$$\begin{aligned} \max & 4x_1 + 2x_2 \\ & 2x_1 \leq 40 \\ & 3x_2 \leq 75 \\ & 3x_1 + 2x_2 \leq 100 \\ & x_1, x_2 \geq 0 \end{aligned}$$

Optimal point?



## To pea or not to pea.

4\$ for peas. 2\$ bushel of carrots.

$x_1$  - to pea!  $x_2$  to carrot ?

Money  $4x_1 + 2x_2$  maximize  $\max 4x_1 + 2x_2$ .

Carrots take 2 unit of water/bushel.

Peas take 3 units of water/bushel. 100 units of water.

$$3x_1 + 2x_2 \leq 100$$

Peas 2 yards/bushel of sunny land. Have 40 sq yards.

$$2x_1 \leq 40$$

Carrots get 3 yards/bushel of shady land. Have 75 sq. yards.


$$3x_2 \leq 75$$

Can't make negative!  $x_1, x_2 \geq 0$ .

A linear program.

$$\begin{aligned} \max & 4x_1 + 2x_2 \\ & 2x_1 \leq 40 \\ & 3x_2 \leq 75 \\ & 3x_1 + 2x_2 \leq 100 \\ & x_1, x_2 \geq 0 \end{aligned}$$

## Feasible Region.




Convex.


Any two points in region connected by a line in region.

Algebraically:

$$\begin{aligned} \text{If } x \text{ and } x' \text{ satisfy onstraint,} \\ \rightarrow x'' = \alpha x + (1 - \alpha)x \end{aligned}$$



Optimal at pointy part of feasible region!  
 Vertex of region.  
 Intersection of two of the constraints (lines in two dimensions)!  
 Try every vertex! Choose best.  
 $O(m^2)$  if  $m$  constraints and 2 variables.  
 For  $n$  variables,  $m$  constraints, how many?  
 $nm$ ?  $\binom{m}{n}$ ?  $n + m$ ?  
 $\binom{m}{n}$   
 Finite!!!!!!  
 Exponential in the number of variables.



$$\begin{aligned} \max & 4x_1 + 2x_2 \\ & 3x_1 \leq 60 \\ & 3x_2 \leq 75 \\ & 3x_1 + 2x_2 \leq 100 \\ & x_1, x_2 \geq 0 \end{aligned}$$

Simplex: Start at vertex. Move to better neighboring vertex.  
 Until no better neighbor. This example.  
 $(0,0)$  objective 0.  $\rightarrow (0,25)$  objective 50.  
 $\rightarrow (16\frac{2}{3}, 25)$  objective  $115\frac{2}{3}$   
 $\rightarrow (20,20)$  objective 120.  
 Duality:  
 Add blue equations to get objective function?  
 $1/3$  times first plus second.  
 Get  $4x_1 + 2x_2 \leq 120$ . Every solution must satisfy this inequality!  
 Objective value: 120.  
 Can we do better? No!  
 Dual problem: add equations to get best upper bound.

## Duality.

$$\begin{aligned} \max & x_1 + 8x_2 \\ & x_1 \leq 4 \\ & x_2 \leq 3 \\ & x_1 + 2x_2 \leq 7 \\ & x_1, x_2 \geq 0 \end{aligned}$$

One Solution:  $x_1 = 1, x_2 = 3$ . Value is 25.  
 Best possible?  
 For any solution.  
 $x_1 \leq 4$  and  $x_2 \leq 3$  ..  
 ....so  $x_1 + 8x_2 \leq 4 + 8(3) = 28$ .  
 Added equation 1 and 8 times equation 2  
 yields bound on objective..  
 Better solution?  
 Better upper bound?

## Duality.

$$\begin{aligned} \max & x_1 + 8x_2 \\ & x_1 \leq 4 \\ & x_2 \leq 3 \\ & x_1 + 2x_2 \leq 7 \\ & x_1, x_2 \geq 0 \end{aligned}$$

Solution value: 25.  
 Add equation 1 and 8 times equation 2 gives..  
 $x_1 + 8x_2 \leq 4 + 24 = 28$ .

Better way to add equations to get bound on function?  
 Sure: 6 times equation 2 and 1 times equation 3.  
 $x_1 + 8x_2 \leq 6(3) + 7 = 25$ .

Thus, the value is at most 25.

The upper bound is same as solution!

Proof of optimality!

## Duality:example

Idea: Add up positive linear combination of inequalities to "get"  
 upper bound on optimization function.

Will this always work?

How to find best upper bound?

## Duality: computing upper bound.

Best Upper Bound.

Multiplier	Inequality
$y_1$	$x_1 \leq 4$
$y_2$	$x_2 \leq 3$
$y_3$	$x_1 + 2x_2 \leq 7$

Adding equations thusly...

$$(y_1 + y_3)x_1 + (y_2 + 2y_3)x_2 \leq 4y_1 + 3y_2 + 7y_3.$$

The left hand side should "dominate" optimization function:

If  $y_1, y_2, y_3 \geq 0$   
 and  $y_1 + y_3 \geq 1$  and  $y_2 + 2y_3 \geq 8$  then..

$$x_1 + 8x_2 \leq 4y_1 + 3y_2 + 7y_3$$

Find best  $y_i$ 's to minimize upper bound?

## The dual, the dual, the dual.

Find best  $y_i$ 's to minimize upper bound?

Again: If you find  $y_1, y_2, y_3 \geq 0$   
and  $y_1 + y_3 \geq 1$  and  $y_2 + 2y_3 \geq 8$  then..  
 $x_1 + 8x_2 \leq 4y_1 + 3y_2 + 7y_3$

$$\begin{aligned} \min & 4y_1 + 3y_2 + 7y_3 \\ & y_1 + y_3 \geq 1 \\ & y_2 + 2y_3 \geq 8 \\ & y_1, y_2, y_3 \geq 0 \end{aligned}$$

A linear program.


The **Dual** linear program.

Primal:  $(x_1, x_2) = (1, 3)$ ; Dual:  $(y_1, y_2, y_3) = (0, 6, 1)$ .

Value of both is 25!

Primal is optimal ... and dual is optimal!

## Again: simplex



$$\begin{aligned} \max & 4x_1 + 2x_2 \\ & 3x_1 \leq 60 \\ & 3x_2 \leq 75 \\ & 3x_1 + 2x_2 \leq 100 \\ & x_1, x_2 \geq 0 \end{aligned}$$

Simplex: Start at vertex. Move to better neighboring vertex.

Until no better neighbor. Duality:

Add blue equations to get objective function?

1/3 times first plus second.

Get  $4x_1 + 2x_2 \leq 120$ . Every solution must satisfy this inequality!

**Geometrically and Complementary slackness:**

Add tight constraints to "dominate objective function."

**Don't add this equation! Shifts.**

## The dual.

In general.

<u>Primal LP</u>	<u>Dual LP</u>
$\max c \cdot x$	$\min y^T b$
$Ax \leq b$	$y^T A \geq c$
$x \geq 0$	$y \geq 0$

**Theorem:** If a linear program has a bounded value, then its dual is bounded and has the same value.

Weak Duality: primal ( $P$ )  $\leq$  dual ( $D$ )

Feasible  $(x, y)$

$$P(x) = c \cdot x \leq y^T Ax \leq y^T b \cdot x = D(y).$$

Strong Duality: next lecture.

## Example: review.

$\max x_1 + 8x_2$	$\min 4y_1 + 3y_2 + 7y_3$
$x_1 \leq 4$	$y_1 + y_3 \geq 1$
$x_2 \leq 3$	$y_2 + 2y_3 \geq 8$
$x_1 + 2x_2 \leq 7$	$x_1, x_2 \geq 0$
$y_1, y_2, y_3 \geq 0$	

"Matrix form"

$\max [1, 8] \cdot [x_1, x_2]$	$\min [4, 3, 7] \cdot [y_1, y_2, y_3]$
$\begin{pmatrix} 1 & 0 \\ 0 & 1 \\ 1 & 2 \end{pmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} \leq \begin{bmatrix} 4 \\ 3 \\ 7 \end{bmatrix}$	$[y_1, y_2, y_3] \begin{pmatrix} 1 & 0 \\ 0 & 1 \\ 1 & 2 \end{pmatrix} \geq \begin{bmatrix} 1 \\ 8 \end{bmatrix}$
$[x_1, x_2] \geq 0$	$[y_1, y_2, y_3] \geq 0$

## Complementary Slackness

<u>Primal LP</u>	<u>Dual LP</u>
$\max c \cdot x$	$\min y^T b$
$Ax \leq b$	$y^T A \geq c$
$x \geq 0$	$y \geq 0$

Given  $A, b, c$ , and feasible solutions  $x$  and  $y$ .

Solutions  $x$  and  $y$  are both optimal if and only if  $x_i(c_i - (y^T A)_i) = 0$ , and  $y_j(b_j - (Ax)_j) = 0$ .

$$x_i(c_i - (y^T A)_i) = 0 \rightarrow$$

$$\sum_i (c_i - (y^T A)_i)x_i = cx - y^T Ax \rightarrow cx = y^T Ax.$$

$$y_j(b_j - (Ax)_j) = 0 \rightarrow$$

$$\sum_j y_j(b_j - (Ax)_j) = yb - y^T Ax \rightarrow by = y^T Ax.$$

$$cx = by.$$

If both are feasible,  $cx \leq by$ , so must be optimal.

In words: nonzero dual variables only for tight constraints!

## Matrix equations.

$\max [1, 8] \cdot [x_1, x_2]$	$\min [4, 3, 7] \cdot [y_1, y_2, y_3]$
$\begin{pmatrix} 1 & 0 \\ 0 & 1 \\ 1 & 2 \end{pmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} \leq \begin{bmatrix} 4 \\ 3 \\ 7 \end{bmatrix}$	$[y_1, y_2, y_3] \begin{pmatrix} 1 & 0 \\ 0 & 1 \\ 1 & 2 \end{pmatrix} \geq \begin{bmatrix} 1 \\ 8 \end{bmatrix}$
$[x_1, x_2] \geq 0$	$[y_1, y_2, y_3] \geq 0$

We can rewrite the above in matrix form.

$$A = \begin{pmatrix} 1 & 0 \\ 0 & 1 \\ 1 & 2 \end{pmatrix} \quad c = [1, 8] \quad b = [4, 3, 7]$$

The primal is  $Ax \leq b, \max c \cdot x, x \geq 0$ .

The dual is  $y^T A \geq c, \min b \cdot y, y \geq 0$ .

### Rules for School...

or..."Rules for taking duals"

Standard:

$$Ax \leq b, \max cx, x \geq 0 \leftrightarrow y^T A \geq c, \min by, y \geq 0.$$

min ↔ max

≥ ↔ ≤

"inequalities" ↔ "nonnegative variables"

"nonnegative variables" ↔ "inequalities"

One more useful trick: Equality constraints.

"equalities" ↔ "unrestricted variables."

### Maximum Weight Matching.

Bipartite Graph  $G = (V, E)$ ,  $w : E \rightarrow Z$ .  
 Find maximum weight perfect matching.  
 Solution:  $x_e$  indicates whether edge  $e$  is in matching.

$$\begin{aligned} &\max \sum_e w_e x_e \\ \forall v : &\sum_{e=(u,v)} x_e = 1 && p_v \\ &x_e \geq 0 \end{aligned}$$

Dual.  
 Variable for each constraint.  $p_v$  unrestricted.  
 Constraint for each variable. Edge  $e$ ,  $p_u + p_v \geq w_e$   
 Objective function from right hand side.  $\min \sum_v p_v$

$$\begin{aligned} &\min \sum_v p_v \\ \forall e = (u, v) : &p_u + p_v \geq w_e \end{aligned}$$

Weak duality? Price function upper bounds matching.

$$\sum_{e \in M} w_e x_e \leq \sum_{e=(u,v) \in M} p_u + p_v \leq \sum_v p_v.$$

Strong Duality? Same value solutions. Hungarian algorithm !!!

### Complementary Slackness.

$$\begin{aligned} &\max \sum_e w_e x_e \\ \forall v : &\sum_{e=(u,v)} x_e = 1 && p_v \\ &x_e \geq 0 \end{aligned}$$

Dual:  
 $\min \sum_v p_v$   
 $\forall e = (u, v) : p_u + p_v \geq w_e$

Complementary slackness:  
 Only match on tight edges.  
 Nonzero  $p_u$  on matched  $u$ .

### Multicommodity Flow.

Given  $G = (V, E)$ , and capacity function  $c : E \rightarrow Z$ , and pairs  $(s_1, t_1), \dots, (s_k, t_k)$  with demands  $d_1, \dots, d_k$ .  
 Route  $D_j$  flow for each  $s_j, t_j$  pair, so every edge has  $\leq \mu c(e)$  flow with minimum  $\mu$ .

variables:  $f_p$  flow on path  $p$ .  
 $P_i$  -set of paths with endpoints  $s_i, t_i$ .

$$\begin{aligned} &\min \mu \\ \forall e : &\sum_{p \ni e} f_p \leq \mu c_e \\ \forall i : &\sum_{p \in P_i} f_p = D_i \\ &f_p \geq 0 \end{aligned}$$

### Take the dual.

$$\begin{aligned} &\min \mu \\ \forall e : &\sum_{p \ni e} f_p \leq \mu c_e \\ \forall i : &\sum_{p \in P_i} f_p = D_i \\ &f_p \geq 0 \end{aligned}$$

Modify to make it  $\geq$ , which "go with min."  
 And only constants on right hand side.

$$\begin{aligned} &\min \mu \\ \forall e : &\mu c_e - \sum_{p \ni e} f_p \geq 0 \\ \forall i : &\sum_{p \in P_i} f_p = D_i \\ &f_p \geq 0 \end{aligned}$$

## Dual.

$$\begin{aligned} \min \mu \\ \forall e: \mu c_e - \sum_{p \ni e} f_p &\geq 0 & d_e \\ \forall i: \sum_{p \in P_i} f_p &= D_i & d_i \\ f_p &\geq 0 \end{aligned}$$

Introduce variable for each constraint.

Introduce constraint for each var:

$$\mu \rightarrow \sum_e c_e d_e = 1. \quad f_p \rightarrow \forall p \in P_i \quad d_i - \sum_{e \in p} d_e \leq 0.$$

Objective: right hand sides.  $\max \sum_i D_i d_i$

$$\begin{aligned} \max \sum_i D_i d_i \\ \forall p \in P_i: d_i \leq \sum_{e \in p} d(e) \end{aligned}$$

$d_i$  - shortest  $s_i, t_i$  path length. Toll problem!

Weak duality: toll lower bounds routing.

Strong Duality. Tight lower bound. First lecture. Or Experts.

Complementary Slackness: only route on shortest paths

only have toll on congested edges.

See you on Thursday.

## Exponential size.

Multicommodity flow.

$$\begin{aligned} \min \mu \\ \forall e: \mu c_e - \sum_{p \ni e} f_p &\geq 0 \\ \forall i: \sum_{p \in P_i} f_p &= d_i \\ f_p &\geq 0 \end{aligned}$$

Dual is.

$$\begin{aligned} \max \sum_i D_i d_i \\ \forall p \in P_i: d_i \leq \sum_{e \in p} d(e) \end{aligned}$$

Exponential sized programs?

Answer 1: We solved anyway!

Answer 2: Ellipsoid algorithm.


Find violated constraint  $\rightarrow$  poly time algorithm.

Answer 3: there is polynomial sized formulation.

Question: what is it?

## Maximum matching and simplex.

$$\begin{aligned} \max x + y + z \\ x \leq 1 \\ x + z \leq 1 \quad a = 1 \\ z + y \leq 1 \quad b = 1 \\ y \leq 1 \\ x \geq 0 \\ y \geq 0 \\ z \geq 0 \end{aligned}$$



[FD: Ellipsoid algorithm for integer knapsack.](#)

Sum:  $x + 2z + y$ .

